aspire

CYBER X

USER MANUAL

EN

DISCLAIMER
Please read this user manual before use. Failure to follow the instructions in this manual may result in fire, electric shock, personal injury or damage to the device or personal property. Death, or property damage.

The manufacturer assumes no responsibility for any damage caused by mishandling that is beyond normal usage defined in this manual.

RECOMMENDED SETUP
We recommend the exclusive use of Aspire original charger, battery, and accessories. Please check the authenticity of the product. As stated, use of chargers, batteries, components, and/or accessories not specifically produced for this device may lead to malfunction, damage to the product and to the cancellation of the warranty.

HANDLING AND USAGE SAFETY
To reduce the risk of fire, shock, or damage to your vape device keep the product and its accessories away from overheated or excessively humid conditions (-5°C / 23°C).

Your device can be damaged if dropped, burned, punctured, crushed, or if it comes in contact with liquid. If you suspect damage to your device or its battery, discontinue use of your device, as it may cause overheating or injury.

To avoid inhaling excess vapour, you either lock or switch off your device when not in use, especially when carrying it in your pocket, bag or purse.

In the case of unusual or abnormal deformation of one of the components of the device or its accessories, stop using it immediately and contact Customer Service.

COIL MAINTENANCE SAFETY
In order not to damage the coil or burn the cotton do not inhale an empty coil. Let the pod/tank stand for 5 minutes after filling so the cotton can be saturated with e-liquid before use.

REPLACEABLE BATTERIES SAFETY
Do not use non-rechargeable batteries. Do not use damaged, modified or counterfeit batteries. Do not leave the battery/ies in the charger when fully recharged. Do not use your vape device while charging.

PRODUCT DISPOSAL SAFETY
Do not discard this product and battery/ies in the environment, and only dispose of these components at declared collection/disposal points.

HEALTH WARNING SAFETY
Do not use the product if you are pregnant, breastfeeding, generally ill, or suffer from respiratory and/or cardiovascular system diseases.

If you experience any side effects while using the product, if the effect(s) continue after you stop using, consult a healthcare professional immediately.

Package Contents:
* 1 Cyber X Device
* 1 Pod 2 ml (1.6 Ω MC)
* 1 Pod 5 ml (1.2 Ω MC)
* 1 Type-C Cable
* 1 Lanyard Loop
* 1 Lanyard

Components View:

Specifications:
Dimensions: 71 x 17.2 x 23.5mm
Capacity: 2ml / 5ml - 2.2 cm
Coil: 1.2Ω (0.8-1.4Ω) coil, non-replaceable
Battery Capacity: Built-in 1000 mAh
Output: 7-9 W Constant output
Charging: Type-C, 2A
Output: 7-9 V Constant output
Activation: Draw Activation
Mode: 8 Puffs Activated
Mode: Button Activation activated

Pod Filling:
1. Pull to remove the pod.
2. Fill the pod from the side of post and side e-liquid .
3. Plug back the silicone seal firmly.
4. Place the pod back.



Remember to clean the condensation between the atomizer and the device.

Instructions:

Device Condition	Status of Indicators & Details
Current Checking	Flash white 5 times
Pod/Coil Communication Error, the Resistance 3 times within 2s	Flash white 3 times
Battery level - 0.1V-3.0V(5%)	On red
Battery level - 3.0+V-4.2V (6%-100%)	On blue
Low Voltage	On green
Overheat Protection	The LED indicator will flash red for 10 times and the device will stop working.
Overheat Protection	The LED indicator will flash white for 10 times and the device will stop working.
Over Voltage Protection	When charging is abnormal, the LED indicator will flash red when charging is not allowed.

Charging:

Maximum charge current	2A
Recommended charge voltage	5V
Battery Full Flashes Green	Fully charged
Overcharge Protection	Protect the Type-C Cable
LED Indicator Flashes green 3 times	Unplug the Type-C Cable

IMPORTANT INFORMATION ABOUT USING NICOTINE-CONTAINING E-LIQUIDS

Before using nicotine containing e-liquid the user should read the following information carefully.

Protection Features:

Over-Time Protection (10s)	The LED indicator will flash red and the device will stop working. Release the fire button to stop.
Short Circuit Protection	The LED indicator will flash red and the device will stop working.
Low Voltage Protection	The LED indicator will flash red.
Overheat Protection	Once the heater is fully charged, the LED indicator will flash green 10 times and the device will stop working.
Overheat Protection	The LED indicator will stay yellow for 10 seconds and the device will stop working.
Over Voltage Protection	When charging is abnormal, the LED indicator will flash red when charging is not allowed.

Charging:

Maximum charge current	2A
Recommended charge voltage	5V
Battery Full Flashes Green	Fully charged
Overcharge Protection	Protect the Type-C Cable

IMPORTANT INFORMATION ABOUT USING NICOTINE-CONTAINING E-LIQUIDS

Before using nicotine containing e-liquid the user should read the following information carefully.

1. NICOTINE CONTAINING E-LIQUID:
Nicotine is a highly addictive substance, it should be noted that nicotine is not only addictive but also toxic. Nicotine, in fact, is more toxic than lead and should be treated with the same precautions as other poisons.

2. NICOTINE WARNINGS AND PRECAUTIONS
* Nicotine is highly addictive and should be treated with caution.
* Harmful if swallowed.
* Do not swallow under the finger cap.
* Those experiencing general ill-health, such as heart problems, should consult their doctor before using nicotine containing products.
* Nicotine may affect the foetus/embryo, so pregnant women should consult their doctor before using nicotine containing products.
* Nicotine may affect breast milk, so breastfeeding women should consult their doctor before using nicotine containing products.
* Nicotine may affect adolescent development, so adolescents should consult their doctor before using nicotine containing products.
* Nicotine may affect the elderly, so the elderly should consult their doctor before using nicotine containing products.
* Nicotine may affect those taking medication, so those taking medication should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from diabetes, so those suffering from diabetes should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from high blood pressure, so those suffering from high blood pressure should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from heart disease, so those suffering from heart disease should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from kidney disease, so those suffering from kidney disease should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from liver disease, so those suffering from liver disease should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from stomach ulcers, so those suffering from stomach ulcers should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from overactive thyroid, so those suffering from overactive thyroid should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from phaeochromocytoma, so those suffering from phaeochromocytoma should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from insulinoma, so those suffering from insulinoma should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from epilepsy, so those suffering from epilepsy should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from migraines, so those suffering from migraines should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from asthma, so those suffering from asthma should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from allergies, so those suffering from allergies should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from eczema, so those suffering from eczema should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from psoriasis, so those suffering from psoriasis should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from lichen planus, so those suffering from lichen planus should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from lupus, so those suffering from lupus should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from scleroderma, so those suffering from scleroderma should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from dermatomyositis, so those suffering from dermatomyositis should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from vitiligo, so those suffering from vitiligo should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from alopecia, so those suffering from alopecia should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from hirsutism, so those suffering from hirsutism should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from hypertrichosis, so those suffering from hypertrichosis should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from hypotrichosis, so those suffering from hypotrichosis should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from anorexia, so those suffering from anorexia should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from bulimia, so those suffering from bulimia should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from obesity, so those suffering from obesity should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from underweight, so those suffering from underweight should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from overweight, so those suffering from overweight should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from malnutrition, so those suffering from malnutrition should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from vitamin deficiency, so those suffering from vitamin deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from mineral deficiency, so those suffering from mineral deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from protein deficiency, so those suffering from protein deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from carbohydrate deficiency, so those suffering from carbohydrate deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from fat deficiency, so those suffering from fat deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from fibre deficiency, so those suffering from fibre deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from water deficiency, so those suffering from water deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from electrolyte deficiency, so those suffering from electrolyte deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from acid-base deficiency, so those suffering from acid-base deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from hormone deficiency, so those suffering from hormone deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from enzyme deficiency, so those suffering from enzyme deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from coenzyme deficiency, so those suffering from coenzyme deficiency should consult their doctor before using nicotine containing products.
* Nicotine may affect those suffering from cofactor deficiency, so those suffering from cofactor deficiency